



Oribatid mite communities on former clay quarries under different reclamation strategy

Todria Nino^a, Murvanidze Maka^a, Mumladze Levan^b

^aInstitute of Entomology, Agricultural University of Georgia. 240, D. Aghmashenebely Alley, Tbilisi, 0156, Georgia

²Institute of Ecology and Institute of Zoology, Ilia State University. 3/5, Cholokashvili Str., Tbilisi, 0165, Georgia

Received: 12 April 2019; accepted: 10 June 2019

A B S T R A C T

In order to establish or regenerate ecosystems in severely disturbed mining sites, reclamation activities are common practice. However, the effectiveness of reclamation strategy might be depended on the specific conditions of the area under question. To evaluate the effect of provided reclamation activities, we investigated structure and composition of oribatid mite communities at an early succession stage in the abandoned Kaspi clay pit. Effectiveness of natural succession and two alternative reclamation activities was evaluated: 1. M1 - do-nothing approach on land under natural secondary succession; 2. M2 - smoothed ground surface with sown seed mixture of herbs and pasture grasses; 3. M3 -three fenced plots with sown grasses and tree seedlings. Neighboring overgrazed natural meadow is chosen as control. Results show that the “do-nothing” approach with leaving post-mining sites without any reclamation is the worse strategy in development of soil oribatid community on lands without “source” area for vegetation (and consequently for soil humus layer) development. Reclaimed fenced areas were relatively poor by oribatid species richness compared to surrounding reclaimed (but not fenced) or natural areas. High abundance of stress tolerant species in fenced sites shows that the time interval between fencing (2012, 2013 and 2014) and sampling (2015-2016) was not enough for soil structure and faunal recovery and longer period is needed to establish sustainable oribatid communities.

Keywords: Oribatid mites, Colonization, Reclamation, Mining, Secondary succession, Overgrazing.

*Corresponding author: Maka Murvanidze; E-mail address: m.murvanidze@agruni.edu.ge

1. Introduction

Natural regeneration of destroyed soil ecosystems (e.g. abandoned open mining areas) could take a long time [1-3]. Reclamation activities are frequently used to enhance ecosystem regeneration processes and help establishment of the biota in disturbed areas. However development of viable soil communities may still need several decades depending on the structure of reclaimed soil [4-6], distance from colonization sources [7-10], age of restored sites [11, 12] etc. Neighboring areas usually serve as a main source for colonization of soil arthropods, however the rate and mode of colonization is supposedly taxon specific. For instance, active dispersal ability of oribatid mites (one of the most abundant and ecologically important groups) in the soil is limited to several centimeters [13]

while ground beetles are fast, long distance runners. It has also been shown that the dispersal rate in slow-moving taxa (such as soil mites) can be assisted by passive pathways so that the species could frequently persist in remote non-viable habitat patches [14] and may play an important (if not concomitant) role in the structuring of mites communities at early stage of succession [13, 15-17]. Accordingly, deep knowledge of the processes that support the development of soil arthropod communities can help establishment in successful reclamation strategies.

Different activities can be employed during the reclamation process including but not limited to re-shaping site surface, planting trees or sowing grasses, applying fertilizers and adding organic products and so on [18]. Depending on the target ecosystem component, different reclamation activities can be used. For instance, regeneration soil and its biota

(as example in an open mining areas with heavily degraded soil surfaces) one has to provide soil substrate and allow recolonization of soil dwelling animals. In addition, other aids can also be applied to support soil process (e.g. sowing grasses). Accordingly, adopting particular reclamation strategy can have a significant role on regeneration efficiency.

Unfortunately there are still insufficient investigations studding soil recovery process in open mining sites and in particular recovery of arthropod communities [19] ensuring no effective reclamation management. Furthermore, any particular reclamation activities undertaken until now in Georgia are only based on expert opinions rather than scientifically informed [20] usually followed no post reclamation monitoring and evaluations.

In this work we investigated the effect of provided reclamation activities on the structure and composition of oribatid mite communities at an early succession stage in the abandoned Kaspi clay pit. Oribatid mites are one of the most diverse and abundant group in soil matrix [21] and play an essential role in soil food web [22]. In addition, diversity of oribatid mites are the most intensively studied in Georgia [23] making this group useful for ecosystem evaluation [24]. Thus, unraveling the effects of reclamation activities on the development of oribatid mite community can help to better understand the succession of soil process. In 2012 Heidelberg Cement Caucasus Company began reclamation of the former clay pit area applying different management strategy including different combinations of surfaceres haping, herb sowing, providing organic source and fencing [20]. Our aim was to evaluate the response of oribatid communities towards provided reclamation activities and to identify the best management strategies. We hypothesized that the fenced and smoothed surface with sowed grass and added organic content supports fastest recolonization and highest diversity of oribatid mites due to higher food supply and less physical disturbance; Though natural succession can lead to formation of diverse oribatid community [10], we were keen to understand how much the community succession rate depends on regeneration time provided in absence of physical disturbance (such as trampling).

2. Material and Methods

2.1. Study area

Kaspi clay pit represents a south western suburb of the Kaspi city (Georgia) (41.93; 44.39). The cli-

mate of the area is sub-Mediterranean with average annual temperature 13°C and annual precipitation 490 mm. The area of the former pit represents a flat depression on the left bank of the river Kura (Fig. 1) with license area of approximately 100 ha and open Kaspi clay pit area - 28.32 ha [20]. After 50th of the last century irregular mining activities were initiated that continued until 2006, after that time Heidelberg Cement Caucasus Ltd resurrected active mining that lasted until 2010. The quarry is surrounded by farmland area, is covered by ruderal vegetation and is extensively used for livestock grazing. For reclamation purposes three type of management have been applied: approach 1. Do-nothing – about 10 ha of area was left untouched with natural secondary succession (M1 – first management category); approach 2 – reclamation activities included smoothed ground surface and sowing commercial seed mixture with herbs and pasture grasses (M2 – second management category); approach 3 – three plots in reclaimed area were fenced each in 2012, 2013 and 2014 to restrict cattle movement. In each plot hay was added immediately as a supply of organic source (M3 – third management category) and seedlings of different species (mostly *Elaeagnus angustifolia* (Russian olive) and *Celtis caucasica* (Caucasian hackberry)) were planted (Table 1; Fig. 1).

Unreclaimed area (M1) is represented with small irregular hills left after mechanical relocation of ground during the mining process (Fig. 1, Table 1). All the areas but M3 were subject of continuous overgrazing.



Fig. 1. Kaspi clay pit with identification of sampling sites. Red solid line indicates area of reclaimed pit (M2), green solid lines show fenced plots (M3) and green dashed line shows area with natural succession (M1).

Table 1. Description of investigation sites with indication of observed and estimated (Chao) species richness. In the rightmost columns, site community data from the same management categories were pooled to evaluate overall management richness.

| Management Category | Sampling units | Site description | Coordinates | Site specific species richness | | Management specific species richness | |
|---------------------|------------------|--|-----------------|--------------------------------|--------------------|--------------------------------------|--------------------|
| | | | | Species Richness | Estimated Richness | Species richness | Estimated richness |
| Ctr | Control | Natural overgrazed meadow. Plant coverage ~ 30% | N41°928 E44°391 | 23 | 30 | 23 | 29 |
| M1 | Not reclaimed | Natural succession on not reclaimed quarry. Overgrazed hills. Plant coverage < 20% | N41°929 E44°393 | 8 | 8 | 8 | 9 |
| M2-1 | Reclaimed | Natural seedlings of <i>Tamarix</i> | N41°927 E44°399 | 17 | 17 | 29 | 32 |
| M2-2 | Reclaimed | Grass sowed in 2012; Overgrazed; Plant coverage ~ 30% | N41°930 E44°527 | 23 | 24 | | |
| M2-3 | Reclaimed | Grass sowed in 2012; Overgrazed; Plant coverage ~ 30% | N41°927 E44°398 | 17 | 19 | | |
| M3-1 | Reclaimed fenced | Fenced in 2014; No grazing; Plant coverage ~ 50% | N41°929 E44°397 | 11 | 22 | 22 | 24 |
| M3-2 | Reclaimed fenced | Fenced in 2013; No grazing; Plant coverage ~ 80% | N41°928 E44°396 | 10 | 10 | | |
| M3-3 | Reclaimed fenced | Fenced in 2012. No grazing; Plant coverage > 90% | N41°930 E44°340 | 15 | 15 | | |

2.2. Sampling

In each management category (M1, M2, M3) and the surrounded control area (that was unaffected at least by mechanical disturbance), four 10 m³ soil samples were taken randomly using soil corer in October 2015, February, May and July of 2016 (covering all four seasons). Soil samples were delivered in the laboratory and invertebrates were extracted using modified Berlese-Tullgren apparatus. Oribatid mites were collected in every 24 hours during one week. Extracted individuals were stored in 70% ethanol; temporary slides were made with lactic acid using hollow ground slides. Only adults were identified to species level using mainly keys of Weigmann [25] and Ghilarov and Krivolutsky [26]. Classification follows Schatz et al [27].

2.3. Data analysis

To compare species diversity and total abundance between different management units, we calculated total species richness for each management category as well as average species richness and individual

density (per square meter) for each sampling unit. Later on, mean values were compared between management units using randomized block ANOVA where sampling period was used as a blocking factor and the management as a treatment. In addition to raw data, nonparametric estimators (Chao indices) and Simpson’s diversity (1 - D) measures were used to consider sampling incompleteness in data analyses [28]. Cluster of faunal similarities we constructed using PAST software and based on Jaccard’s similarity index.

3. Results

In total 3055 oribatid individuals belonging to 43 species from 25 families were identified (Table 2). Highest total number of species (29) was found in reclaimed overgrazed areas (M2-1 – M2-3) while lowest number (8 species) in not-reclaimed site (M1). Accordingly, this site showed lowest index of diversity (1-D = 0.15) (Table 1 and 3). Non-parametric estimators (Chao 1) showed the same trend and indicated a near complete inventory of species in almost all cases except Ctr where significantly

more species are expected than sampled (Table 1). 82% of individuals were represented by 7 dominant species only (*Punctoribates punctum* (C.L. Koch, 1839), *Oribatula* (*Z.*) *cognata* (Oudemans, 1902), *Pilogalumna crassiclava* (Berlese, 1914), *Tectocephus velatus* (Michael, 1880), *Epilohmannia cylindrica* (Berlese, 1904), *Acrotrititia ardua* (C.L. Koch,

1841), *Ramusella clavipectinata* (Michael, 1885)) among which *P. punctum* composed 56% of total individuals and dominated in all samples. Rare species *Scutovertex armazi* Murvanidze & Weigmann, 2012, was found on natural meadow. Up to date this species was known from type locality only (Armazi Mountain) [29].

Table 2. List of Oribatid mites found on Kaspi clay pit with number of individuals for each site

| Species | M1 | M2-1 | M2-2 | M2-3 | M3-1 | M3-2 | M3-3 | CTR |
|---|-----|------|------|------|------|------|------|-----|
| <i>Epilohmannia cylindrica</i> (Berlese, 1904) | 3 | 8 | 7 | 49 | 1 | 14 | 68 | 14 |
| <i>Acrotrititia ardua</i> (C.L. Koch, 1841) | 13 | 7 | 18 | 16 | 0 | 9 | 24 | 12 |
| <i>Nothrus anauniensis</i> (Canestrini & Fanzago 1876) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 4 |
| <i>Trhypochthonius tectorum</i> (Berlese, 1896) | 0 | 2 | 0 | 0 | 0 | 0 | 4 | 0 |
| <i>Belba dubinini</i> Bulanova (Zachvatkina, 1962) | 0 | 1 | 17 | 6 | 0 | 7 | 3 | 1 |
| <i>Metabelba italica</i> (Sellnick, 1931) | 0 | 0 | 0 | 2 | 0 | 0 | 0 | 0 |
| <i>Berlesezetes cuspidatus</i> (Mahunka, 1982) | 0 | 0 | 0 | 0 | 2 | 0 | 0 | 0 |
| <i>Dorycranosus splendens</i> (Coggi, 1898) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 |
| <i>Liacarus brevilamellatus</i> (Mihelcic, 1955) | 0 | 0 | 0 | 0 | 0 | 0 | 8 | 0 |
| <i>Xenillus tegeocranus</i> (Hermann, 1804) | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 2 |
| <i>Ceratoppia quadridentata</i> (Haller, 1882) | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 |
| <i>Carabodes</i> sp. | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 |
| <i>Epimerella smirnovi</i> (Kulijev, 1962) | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 |
| <i>Oppiella nova</i> (Oudemans, 1902) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2 |
| <i>O. fallax</i> (Paoli, 1908) | 0 | 0 | 0 | 3 | 0 | 0 | 0 | 0 |
| <i>O. subpectinata</i> (Oudemans, 1900) | 0 | 0 | 0 | 2 | 0 | 0 | 0 | 0 |
| <i>Ramusella clavipectinata</i> (Michael 1885) | 1 | 65 | 7 | 50 | 1 | 2 | 13 | 5 |
| <i>Quadroppia quadricarinata</i> (Michael 1885) | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 |
| <i>Suctobelbella subtrigona</i> (Oudemans, 1916) | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 |
| <i>Tectocephus velatus velatus</i> (Michael, 1880) | 0 | 13 | 28 | 10 | 0 | 5 | 12 | 2 |
| <i>T. velatus sarekensis</i> (Trägårdh, 1910) | 6 | 13 | 0 | 12 | 1 | 0 | 9 | 0 |
| <i>Tectoribates ornatus</i> (Schuster, 1958) | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 1 |
| <i>Camisia horrida</i> (Hermann 1804) | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| <i>Scutovertex armazi</i> (Murvanidze & Weigmann, 2012) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 |
| <i>S. minutes</i> (Koch, 1836) | 0 | 2 | 2 | 6 | 0 | 0 | 8 | 4 |
| <i>S. sculptus</i> (Michael, 1879) | 0 | 4 | 3 | 5 | 1 | 1 | 0 | 7 |
| <i>Peloptulus phaenotus</i> (C.L. Koch, 1844) | 0 | 0 | 0 | 0 | 0 | 0 | 3 | 0 |
| <i>Parachipteria fanzagoi</i> (Jacot, 1929) | 0 | 0 | 0 | 2 | 0 | 0 | 0 | 0 |
| <i>Oribatella colchica</i> (Krivolutsky, 1974) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 |
| <i>Oribatula tibialis</i> (Nicolet, 1855) | 0 | 0 | 0 | 0 | 3 | 0 | 0 | 0 |
| <i>O. (Z.) cognata</i> (Oudemans 1902) | 2 | 1 | 43 | 97 | 12 | 0 | 12 | 70 |
| <i>O. (Z.) exavata</i> (Berlese, 1916) | 0 | 3 | 6 | 0 | 0 | 0 | 0 | 0 |
| <i>O. (Z.) frisiae</i> (Oudemans, 1900) | 2 | 38 | 0 | 0 | 0 | 0 | 0 | 0 |
| <i>O. (Z.) terricola</i> (V.D. Hammen, 1952) | 0 | 8 | 13 | 0 | 0 | 0 | 0 | 0 |
| <i>Simkinia tianschanica</i> (Krivolutsky 1967) | 0 | 0 | 1 | 2 | 0 | 0 | 10 | 32 |
| <i>Schelorbates leavigatus</i> (C.L. Koch 1836) | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 |
| <i>Ceratozetes minutissimus</i> (Willmann, 1951) | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Punctoribates punctum</i> (Koch, 1839) | 335 | 152 | 141 | 247 | 29 | 94 | 450 | 276 |
| <i>Galumna flagellate</i> (Willmann, 1925) | 0 | 0 | 0 | 0 | 0 | 7 | 0 | 107 |
| <i>Pergalumna nervosa</i> (Berlese, 1914) | 0 | 14 | 5 | 2 | 0 | 2 | 5 | 3 |
| <i>Pilogalumna crassiclava</i> (Berlese, 1914) | 0 | 20 | 32 | 34 | 0 | 6 | 6 | 24 |
| <i>P. tenuiclava</i> (Berlese, 1908) | 0 | 0 | 0 | 2 | 0 | 0 | 0 | 0 |
| <i>Protorbates capucinus</i> (Berlese, 1908) | 0 | 12 | 15 | 24 | 1 | 0 | 0 | 1 |

Table 3. Diversity indexes of oribatid mites on former Kaspi clay pit

| | CTR | M1 | M2-1 | M2-2 | M2-3 | M3-1 | M3-2 | M3-3 |
|----------------|-------|-------|-------|-------|-------|-------|-------|-------|
| Taxa_S | 23 | 8 | 17 | 23 | 17 | 11 | 10 | 15 |
| Individuals | 572 | 363 | 363 | 575 | 340 | 53 | 147 | 635 |
| Dominance_D | 0.289 | 0.853 | 0.228 | 0.235 | 0.214 | 0.358 | 0.430 | 0.517 |
| Simpson_1-D | 0.711 | 0.147 | 0.772 | 0.765 | 0.786 | 0.642 | 0.571 | 0.483 |
| Evenness_e^H/S | 0.245 | 0.185 | 0.421 | 0.309 | 0.452 | 0.398 | 0.393 | 0.230 |





Fig. 2. Average abundance of oribatid mites on investigated sites. Error bars indicates ± 1 standard deviation.

ANOVA model describing abundance and richness differences among management classes were significant (for main effects $p(F_{3,91}) = 0.008$ and $p(F_{3,91}) = 0.002$ respectively). However, in case of abundance, only Ctr samples was significantly abundant ($p = 0.027$) compared to M3 samples, while there were no significant differences in other cases

(Fig. 2). Similarly, sample species richness was significantly higher in Ctr sites compared to M3 ($p = 0.006$) and in M2 compared to M3 ($p = 0.046$).

Oribatid faunal composition showed higher similarity between Control and M3 sites while oribatids from unmanaged (M1) site were isolated from those with different management approaches (Fig. 3).

Fig. 3. Cluster of faunal similarity between plots with different management activities (faunal similarity is based on Jaccard's coefficient).



4. Discussion

Mining on Kaspi clay pit was stopped in 2009, while restoration activities were initiated in 2012 [20]. Our results show that the “do-nothing” approach with leaving post-mining sites without any reclamation is not effective strategy in development of soil oribatid community (and most probably whole soil ecosystem). This contradicts with our earlier finding which showed that natural succession can support even more diverse fauna than reclamation [10]. Effectiveness of natural succession is supported by other investigations as well [11, 25, 32]. Poor development of oribatid communities on M1 sites can be explained by two reasons: (1) time interval between stopping mining activities (2009) and sampling (2015) was not enough for development of oribatid fauna. More than 10 years are needed for faunal recovery [10, 19]; (2) there is no “source” area for vegetation (and consequently for soil humus layer) development around the whole area - M1 plots are surrounded by extensively overgrazed natural and reclaimed meadows without any forested territory. Intensive grazing leaves soil bare and exposed to wind erosion which makes it infer-

tile and unsuitable for development of soil fauna. In our previous investigation neighboring forests served as a source for oribatid colonization on dump and abandoned quarry sites [10], hence we can conclude that presence of diverse habitat fragments speeds up secondary succession on former mining sites, while in their absence succession can be either very slow or impossible and active reclamation is strongly suggested.

In spite of high faunal similarity between control and fenced (M3) sites (Figure 2), opposite to our expectations, reclaimed fenced areas (that are protected from grazing) were relatively poor by oribatid species richness (plot species richness as well as total species diversity) compared to surrounding reclaimed (but not fenced) or natural areas. Fencing protects reclaimed sites from heavy overgrazing and leads to formation of dense vegetation cover with favorable conditions for soil fauna. However, as table 3 shows, evenness of species distribution in fenced sites tends to be lower compared to reclaimed unfenced areas and high abundance was provided by dominance of single species – *P. punctum*. This is true for M1 site as well, where high overall abundance is provided by *P. punctum*, when

the rest seven species are presented by minor quantities (Table 2). This species (with another co-dominant species like *T. velatus*) is known as cosmopolite one and an effective colonizer of disturbed habitats in early stages of soil recovery [9, 10, 32]. High abundance of these stress tolerant species in protected sites shows that the time interval between fencing (2012, 2013 and 2014) and sampling (2015-2016) was not enough for soil structure and faunal recovery. Moreover, for sensitive species that are heavily dependent on habitat quality and passive dispersal opportunities, fencing may pose an additional barrier for animals (e.g. cattle) who could contribute to passive dispersal of oribatids. Indeed, occurrence of species such as *Ceratoppia quadridentata* (Haller, 1882), *Quadroppia quadricarinata* (Michael, 1885), *Epimerella smirnovi* (Kulijev, 1962), *S. armazi*, *Oribatella colchica* Krivolutsky, 1974, *Galumna flagellata* Willmann, 1925 in surrounding reclaimed areas which are not fenced, indicates that these species were not able to reach the fenced habitats. Previous investigations show that soil arthropods (and oribatid mites among them) can be passively dispersed by other vectors such as wind [13], birds [16], beetles [32, 34] and even frogs [35]. Nonetheless, it seems that none of above mentioned vector can be only responsible for mite colonization in Kaspi quarry areas. We suppose that the active livestock grazing in postmining sites while hindering effective development of vegetation cover and accumulation of organic material in soil [36], can support active translocation of soil arthropods from surrounding areas. Unfortunately, relatively little is known on the role of livestock in oribatid mite dispersal and our conclusions needs to be further corroborated.

Acknowledgement

This work was supported by Shota Rustaveli National Science Foundation (SRNSF) № PhDF17-124 “Oribatid mites – indicators of the disturbed ecosystems” and Heidelberg Cement company.

References

- [1] P. Andrés, E. Mateos, Soil mesofaunal responses to post-mining restoration treatments, *Appl. S. Ecol.* 33 (2006) 67-78.
- [2] H.B. Redi, R.J. van Aarde, T.D. Wassenaar, 2011. Coastal Dune Forest Development and the Regeneration of Millipede Communities, *Restor. Ecol.* 13 (2) (2011) 284–291.
- [3] K. Prach, K. Rehoukova, J. Rehounek, P. Konvalikova, Ecological Restoration of Central European Mining Sites: A Summary of a Multi-site Analysis, *Landsc. Res.* 36 (2) (2011) 263–268.
- [4] A.S. Zaitsev, N.M. van Straalen, Species diversity and metal accumulation in oribatid mites (Acari, Oribatida) of forests affected by a metallurgical plant. *Pedobiologia* 45 (2001) 467–479.
- [5] P. Skubała, M. Ciosek, Oribatid mites (Acari, Oribatida) colonizing the zinc metallurgic dump, *Fragm. Faun.* 42 (6) (1999), 41-55.
- [6] P. Skubała, K. Rola, P. Osyczka, Oribatid communities and heavy metal bioaccumulation in selected species associated with lichen in a heavily contaminated habitat. *Environ. Sci. Pollut. Res.* 23 (2016) 8861–8871. DOI 10.1007/s11356-016-6100-z.
- [7] P. Skubała, Colonization of a dolomitic dump by oribatid mites (Acari, Oribatida), *Pedobiologia* 43 (1999) 145-159.
- [8] P. Skubała, A. Kafel, Oribatid mite communities and metal bioaccumulation in oribatid species (Acari, Oribatida) along the heavy metal gradient in forest ecosystems, *Environ. Pollut.* 132 (2004) 51-60.
- [9] P. Skubała, M. Gulvik, Pioneer oribatid mite communities (Acari, Oribatida) in newly exposed natural (glacier foreland) and anthropogenic (post-industrial dump) habitats, *Pol. J. Ecol.* 53 (3) (2005) 395–407.
- [10] M. Murvanidze, L. Mumladze, T. Arabuli, Er. Kvavadze, Oribatid mite colonization on sand and manganese tailing sites, *Acarologia* 53 (2) (2013) 127-139.
- [11] W. Dunger, M. Wanner, H. Hauser, K. Hohberg, H.J. Schulz, T. Schwalbe, B. Seifert, J. Vogel, K. Voigtländer, B. Zimdars, K.P. Zülka, Development of soil fauna at mine sites during 46 years after afforestation. *Pedobiologia* 45 (2001) 243-271.
- [12] B.N. McAdams, S.A. Quideau, M.J.B. Swallow, L.M. Lumley, Oribatid mite recovery along a chronosequence of afforested boreal sites following oil sands mining, *Forest Ecol. Manag.* 422 (2018) 281–293.
- [13] R. Lehmitz, D. Russell, K. Hohberg, A. Christian, W.E.R. Xylander, Active dispersal of oribatid mites into young soils, *Appl. Soil Ecol.* 55 (2012) 10–19.
- [14] M. Ingimarsdóttir, T. Caruso, J. Ripa, Ó. B.

- Magnúsdóttir, M. Migliorini, K. Hedlund, Primary assembly of soil communities: disentangling the effect of dispersal and local environment, *Oecologia*, 170 (2012) 745-754. DOI 10.1007/s00442-012-2334-8
- [15] Sh. Karasawa, K. Gotoh, T. Sasaki, N. Hijii, Wind-based dispersal of oribatid mites (Acari: Oribatida) in a subtropical forest in Japan, *J. Acarol. Soc. Jpn*, 14 (2) (2005) 117-122.
- [16] N.V. Lebedeva, Oribatid mites transported by birds to polar islands, a review, In: G. Hempel, K. Lochte, G. Matishov (Eds). *Arctic Marine Biology. Reports on polar and marine research*, Bremerhaven, Alfred Wegener Institute for Polar and Marine Research, 640, 152-161, 2012.
- [17] R. Lehmitz, D. Russell, K. Hohberg, A. Christian, W.E.R. Xylander, Wind dispersal of oribatid mites as a mode of migration, *Pedobiologia* 54 (2011) 201–207.
- [18] R. Tropek, T. Kadlec, M. Hejda, P. Kocarec, J. Skuhrovec, I. Malenovsky, S. Vodka, L. Spitzer, P. Banar, M. Konvicka, Technical reclamations are wasting the conservation potential of post-mining sites. A case study of black coal spoil dumps. *Ecol. Eng.* 43 (2012) 13-18.
- [19] P. Skubała, Do we really need land reclamation on dumps? (Oribatid fauna case studies), In: G. Gabrys, S. Ignatovicz, (Eds), *Advances in Polish Acarology*, Wydawnictwo SGGW, Warszawa, 366-374, 2006.
- [20] U. Tranke, *Restoration concept Kaspi clay pit HeidelbergCement Caucasus ltd*, Blaubeuren, 2012.
- [21] M. Maraun, H. Schatz, S. Sheu, Awesome or ordinary? Global diversity patterns of oribatid mites, *Ecography* (2007) 209-2016.
- [22] K. Schneider, C. Renker, S. Scheu, M. Maraun, Feeding biology of oribatid mites: a minireview, *Phytophaga* (2004), 247 -256.
- [23] M. Murvanidze, L. Mumladze, Annotated checklist of Georgian oribatid mites. *Zootaxa* 4089 (1) (2016) 1-81.
- [24] V. Gergocs, L. Hufnagel, Application of oribatid mites as indicators (Review), *Appl. Ecol. Env. Res.* 7 (1) (2009) 79-98.
- [25] G. Weigmann, *Hornmilben (Oribatida)*, Die Tierwelt Deutschlands, Teil 76, Goecke & Evers. Keltern, 2006.
- [26] M.S. Ghilarov, D.A. Krivolutsky, Identification keys of soil inhabiting mites, Nauka, Moscow, 1975. (in Russian)
- [27] H. Schatz, V.M. Behan-Pelletier, B.M.O Connor, R.A. Norton, Suborder Oribatida van der Hammen, 1968. In: Zhang, Z.Q. (Ed), *Animal biodiversity: An outline of higher level classification and survey of taxonomic richness*. *Zootaxa* 3148 (2011) 141-148.
- [28] A. Magurran, *Measuring Biological Diversity*, Blackwell Publishing, Oxford, RU, 2004.
- [29] M. Murvanidze, G. Weigmann, Two new species of oribatid mites (Acari, Oribatida) *Haplozetes longisacculus* and *Scutovertex armazi* from Georgia (Caucasus), *Acarina* 20 (2) (2012) 167-172.
- [30] P. Skubała, Moss mites (Aearina : Oribatida) on industrial dumps of different ages, *Pedobiologia* 39 (1995)170-184.
- [31] M. Murvanidze, N. Todria, Oribatida diversity on limestone and clay quarries. *Proc. Inst. Zool*, 24 (2015) 159-169.
- [32] W. Dunger, M Wanner, H. Hauser, K. Hohberg, H.J. Schulz, T. Schwalbe, B. Seifert, J. Vogel, K. Voigtländer, B. Zimdars, K.P. Zulka, Development of soil fauna at mine sites during 46 years after afforestation. *Pedobiologia* 45 (2001) 243-271.
- [33] S. Ermilov, A. Frolov, *Ramusella (Dosangoppia) bochkovi*(Acari, Oribatida, Oppiidae), a new subgenus and species of oribatid mites phoretic on *Ceratophyus polyceros* (Pallas, 1771) (Coleoptera, Geotrupidae) from Russia, *Syst. Appl. Acar.* 24 (2) (2018) 209–221.
- [34] E. Waleckx, T. J. Montalvo-Balam, A. Pinzón-Canua, A. Arnal, G. Marti, P.A. Martíneć, First report of phoresy by an oribatid mite (Acari: Oribatida) on a triatomine bug (Hemiptera: Reduviida), *Int. J. Acar.* 44 (4-5) (2018) 210-211. DOI:10.1080/01647954.2018.1487467.
- [35] L.E. Beaty, H.J. Esser, R.M. R.A. Norton, First report of phoresy by an oribatid mite (Trhypochthoniidae: *Archegozetes magnus*) on a frog (Leptodactylidae: *Engystomopspus tulosus*). *Int. J. Acar.* 39 (2013), 325-326.
- [36] P.C.F. Carvalho, I.A.A. de Moraes, E.D. de Souza, R.M. Sulc, C.R. Lang, J.P.C. Flores, M.L. Terra Lopes, J.L.S. da Silva, O. Conte, C. de Lima Wesp, R.L.R.S. Fontaneli, C. Bayer, Managing grazing animals to achieve nutrient cycling and soil improvement in no-till integrated systems. *Nutr.Cycl. Agroecosyst*, 88 (2010) 259–273. DOI 10.1007/s10705-010-9360-x.